

Comparison of Non-human Primate versus Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes for Treatment of Myocardial Infarction.

Journal: Stem Cell Reports

Publication Year: 2018

Authors: Xin Zhao, Haodong Chen, Dan Xiao, Huaxiao Yang, Ilanit Itzhaki, Xulei Qin, Tony Chour, Aitor Aguirre, Kim Lehmann, Youngkyun Kim, Praveen Shukla, Alexandra Holmstrom, Joe Z Zhang, Yan Zhuge, Babacar C Ndoye, Mingtao Zhao, Evgenios Neofytou, Wolfram-Hubertus Zimmermann, Mohit Jain, Joseph C Wu

PubMed link: 29398480

Funding Grants: Macaca mulatta as advanced model for predictive preclinical testing of engineered cardiac autografts and allografts

Public Summary:

Non-human primates (NHPs) can serve as a human-like model to study cell therapy using induced pluripotent stem cell-derived cardiomyocytes (iPSC-CMs). However, whether the efficacy of NHP and human iPSC-CMs is mechanistically similar remains unknown. To examine this, RNU rats received intramyocardial injection of 1×10^7 NHP or human iPSC-CMs or the same number of respective fibroblasts or PBS control ($n = 9-14/\text{group}$) at 4 days after 60-min coronary artery occlusion-reperfusion. Cardiac function and left ventricular remodeling were similarly improved in both iPSC-CM-treated groups. To mimic the ischemic environment in the infarcted heart, both cultured NHP and human iPSC-CMs underwent 24-hr hypoxia in vitro. Both cells and media were collected, and similarities in transcriptomic as well as metabolomic profiles were noted between both groups. In conclusion, both NHP and human iPSC-CMs confer similar cardioprotection in a rodent myocardial infarction model through relatively similar mechanisms via promotion of cell survival, angiogenesis, and inhibition of hypertrophy and fibrosis.

Scientific Abstract:

Non-human primates (NHPs) can serve as a human-like model to study cell therapy using induced pluripotent stem cell-derived cardiomyocytes (iPSC-CMs). However, whether the efficacy of NHP and human iPSC-CMs is mechanistically similar remains unknown. To examine this, RNU rats received intramyocardial injection of 1×10^7 NHP or human iPSC-CMs or the same number of respective fibroblasts or PBS control ($n = 9-14/\text{group}$) at 4 days after 60-min coronary artery occlusion-reperfusion. Cardiac function and left ventricular remodeling were similarly improved in both iPSC-CM-treated groups. To mimic the ischemic environment in the infarcted heart, both cultured NHP and human iPSC-CMs underwent 24-hr hypoxia in vitro. Both cells and media were collected, and similarities in transcriptomic as well as metabolomic profiles were noted between both groups. In conclusion, both NHP and human iPSC-CMs confer similar cardioprotection in a rodent myocardial infarction model through relatively similar mechanisms via promotion of cell survival, angiogenesis, and inhibition of hypertrophy and fibrosis.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/comparison-non-human-primate-versus-human-induced-pluripotent-stem-cell>